



The *In Vitro* Impacts of Some Plant Extracts on Carbonic Anhydrase I, II and Paraoxonase-1

Karbonik Anhidraz I, II ve Paraoksonaz-1 Üzerine Bazı Bitki Ekstraktlarının *In Vitro* Etkileri

Elif Duygu Kaya¹, Bülent Erğün², Yeliz Demir³, Zuhâl Alım⁴, Şükrü Beydemir^{5*}

¹Department of Food Engineering, Faculty of Engineering, Iğdır University, Iğdır, Turkey.

²Department of Pharmaceutical Toxicology, Faculty of Pharmacy, Anadolu University, Eskişehir, Turkey.

³Department of Chemistry, Faculty of Sciences, Ataturk University, Erzurum, Turkey.

⁴Department of Chemistry, Faculty of Science and Arts, Ahi Evran University, Kırşehir, Turkey.

⁵Department of Biochemistry, Faculty of Pharmacy, Anadolu University, Eskişehir, Turkey.

ABSTRACT

The presented article focuses on the *in vitro* inhibition of plant extracts on the human carbonic anhydrase isoforms (hCA I and hCAII), and paraoxonase-1 (PON1) activities. Five different plants (*Alcea rosea*, *Foeniculum vulgare*, *Elettaria cardamomum*, *Laurus azorica* and *Lavandula stoechas*) were selected in this study. Methanol, ethanol, and water extracts of plants were prepared and the concentration-dependent inhibition degrees were found for hCA I and hCA II isozymes and hPON1 spectrophotometrically. Thus, IC_{50} (mg/mL) values were obtained for each extract. Methanolic extract of *Elettaria cardamomum* has the highest inhibitory effects (0.032 mg/mL). The water extracts of plants showed lower inhibitory impacts compared to the methanol and ethanol extracts.

Keywords

Carbonic anhydrase, inhibition, paraoxonase, plant extract.

Öz

Bu makale insan karbonik anhidraz (CA) izoformları hCAI, hCAII ve paraoksonaz-1 (PON1) aktiviteleri üzerine bitki özütlerinin *in vitro* inhibisyonlarına odaklanmaktadır. Bu çalışmada beş farklı bitki (*Alcea rosea*, *Foeniculum vulgare*, *Elettaria cardamomum*, *Laurus azorica* and *Lavandula stoechas*) seçildi. Bitkilerin metanol, etanol ve su özütleri hazırlandı ve hCA I, II izoenzimleri ve hPON1 için derişime bağılı inhibisyon dereceleri spektrofotometrik olarak bulundu. Böylece her bir özüt için IC_{50} (mg/mL) değerleri belirlendi. *Elettaria cardamomum*'un metanol özütü en yüksek inhibisyon etkisine (0.032 mg/mL) sahipti. Bitkilerin su özütleri metanol ve etanol ekstraktları ile kıyaslandığında daha düşük inhibitör etkisi gösterdi.

Anahtar Kelimeler

Karbonik anhidraz, inhibisyon, paraoksonaz, bitki özütü.

Article History: Received: Aug 08, 2018; Revised: Dec 10, 2018; Accepted: Jan 08, 2019; Available Online: Mar 01, 2019.

DOI: 10.15671/HJBC.2019.274

Correspondence to: Ş. Beydemir, Department of Biochemistry, Faculty of Pharmacy, Anadolu University, Eskişehir, Turkey.

E-Mail: sukurbeydemir@anadolu.edu.tr

INTRODUCTION

Plant extracts have been applied for the treatment of many diseases in folk medicine such as diabetes [1] epilepsy [2], Alzheimer [3] and cardiovascular disease [4]. Besides the positive effects of these applications, the number of negative effects is not to be underestimated. Especially, plants are rich in terms of antioxidant components. Thus, they are widely consumed both as fresh food and extract on the world. Antioxidants are used to minimize the many diseases including cancer. Indeed, the source of many metabolic disorders is oxidative changes in the metabolism. The changes may have negative effects on the quality of life. Antioxidants are well known to role as preservatives against to oxidative stress. However, plants include various natural compounds in their structure. These compounds may be important for some disorders of living things. A lot of natural compounds may have still not been purified from plants. Therefore, use of the plant extracts has an important place as preservative or therapeutic in herbal medicine [5].

In the present study, *Alcea rosea* (*A. rosea*), *Foeniculum vulgare* Mill., *Cardamom* (*Elettaria cardamomum*), *Laurus azorica* L., *Lavandula stoechas* were selected as the source of the plant extract. *Alcea rosea* (*A. rosea*) (Family Malvaceae) commonly known as hollyhock which is a traditional Chinese herb [6]. The medicinal parts of *A. rosea* contain seeds, roots and flowers [7]. *A. rosea* flowers and roots have employed as febrifuge, diuretic, expectorant, demulcent, coolant, anti-inflammatory, and astringent agent [8]. *Foeniculum vulgare* Mill. (fennel) is a popular green aromatic herb of the family Apiaceae that is common throughout the Mediterranean region [9]. The essential oil of *Foeniculum vulgare* Mill. seeds (fennel oil) has been used to treat conditions of the reproductive, endocrine respiratory and digestive systems in traditional medicine [10]. Cardamom (*Elettaria cardamomum*), commonly known as the “queen of spices” is one of the highly priced exotic spices in the world. Cardamom is mainly used for domestic culinary purposes and also has medicinal applications as a stimulant, digestive, breath freshener, carminative, and as an aphrodisiac. Cardamom has prominent antioxidant [11], anti-inflammatory [12], and antimicrobial properties [13]. *Laurus azorica* L., commonly known as bay leaves, belongs to Laureacea family, which *Laurus nobilis*. It is widely used as a spicy fragrance and flavor in traditional meat dishes, stews and rice [14]. *Lavandula stoechas* is a species of aromatic flowering plant of the Lamiaceae family [15]. It is used

as cooking spices and fragrance, and its essential oil (EO) is found in the production of food, drinks, perfumes, cosmetics and pharmaceuticals [16-19].

It is well known that enzymes are crucial biocatalyzers in the metabolism. Therefore, all substances taken in the body may interact with various enzymes. Especially some enzymes are called as drug-target and chemical target. Carbonic anhydrase and paraoxonase are known as drug-target enzymes in the literature. Carbonic anhydrases (CA, carbonate hydrolyase, E.C.4.2.1.1) are one of the most studied enzymes [20,21]. They are presented in all species and constitute a family involved in regulating pH and water, electrolyte, and ion transport [22]. The zinc-metalloenzymes play an important role in respiration, carbon dioxide and bicarbonate transport, pH and CO₂ balance in the lungs and tissues, the release of secretions from various tissues and organs, and especially in biosynthetic metabolic processes such as glycogenesis and lipogenesis in mammals [23]. Carbonic anhydrases are also necessary for CO₂ fixation in plants, algae, and prokaryotes. While the CA enzyme is common in organisms, its currently 16 isozymes are known in various tissues [24]. CA catalyzes the converting reaction from CO₂ to HCO₃⁻ which is named hydratase activity. It is a physiological activity of the CA.

On the other hand, CA enzyme exhibits also esterase activity under in vitro conditions. Inhibition or activation of CAs may important in terms of treatment many diseases such as diabetes, cancer, epilepsy, Alzheimer and cardiovascular disease. From this perspective, carbonic anhydrase inhibitors are clinically highly important compounds [25,26].

Paraoxonase (PON) is a mammalian lactonase that especially exists in the liver and HDL-bound enzyme. It is also known as drug target enzyme in the metabolism. Human PON1 (hPON1) is calcium-dependent enzyme. The enzyme consists two calcium ions its three-dimension structure. While one calcium ion is bound the active site, another is on structural part of the enzyme. PON1 consists of 355 amino acids having at least two N-linked carbohydrate chains. PON1 is synthesized in the liver and released from here into the blood [27]. Decrease of PON1 activities may related with some metabolic or genetic disorders such as arthritis, diabetes mellitus, rheumatoid cardiovascular diseases, age-related macular degeneration, chronic renal failure, and hyperthyroidism [27]. Moreover, PON1 has a significant place in HDL metabolism and in the prevention

of atherosclerosis. Some experimental data show that PON1 makes a central contribution to the antioxidant capacity of HDL Paraoxonase, by this antioxidant property prevent from oxidation of both HDL and LDL. Thus, there is a close physiological relation between PON1 and HDL in plasma [28]. HDL facilitates the secretion of the PON1 from the liver, stabilizes the enzyme and provides the hydrophobic environment that is needed for the function of PON1 [29]. It is now known that the effects of chemicals on enzymatic mechanisms are important. Because, almost all reactions in the metabolism are catalyzed by various enzymes. These dramatic changes on the enzymatic mechanisms may lead to many disorders as mentioned above. Therefore, to determine the impacts on enzyme activities of herbal extracts commonly used in alternative medicine is crucial.

The present article focuses on in vitro inhibitory effects on human PON1 and CA I and II isoforms of *Alcea rosea*, *Foeniculum vulgare*, *Elettaria cardamomum*, *Laurus azorica* and *Lavandula stoechas*.

MATERIALS and METHODS

Materials

Paraoxon, p-Nitrophenylacetate, Cyanogen bromide-activated-Sepharose 4B, protein assay reagents and chemicals for electrophoresis were obtained from Sigma-Aldrich Co. (Sigma- Aldrich Chemie GmbH Export Department Eschenstrasse 5, 82024 Taufkirchen, Germany). All other chemicals were analytical grade and obtained from Merck (Merck KGaA Frankfurter Strasse 250, D 64293 Darmstadt, Germany).

Plant material and Extraction

Plants were obtained from a local market at Erzurum, Turkey. Extraction process was done according to previous study [30].

Purification of Carbonic Anhydrase Isozymes and Esterase Activity Assay

CA isoforms were purified using Sepharose-4B-L-tyrosine-sulfanilamide affinity chromatography in a single step [31]. The affinity material, Sepharose-4B-L-tyrosine-sulfanilamide, was prepared conforming to our previous technique [31]. The protein eluates were measured at 280 nm as described previously, spectrophotometrically [32,33]. The activities of CA isoforms were evaluated according to the Verpoorte et al. (1967) [34]. p-Nitrophenylacetate was used as a substrate converted by both isoforms to the

p-nitrophenolate ion in this technique. The absorbance changes were determined during 3 min at 25°C.

Protein Quantity Assay

Bradford procedure was used for determination of protein amount [35]. The bovine serum albumin was used as standard for this evaluation, which was done at 595 nm according to previous studies [36-38].

SDS-Polyacrylamide Gel Electrophoresis

According to Laemmli's procedure (1970) the purity and presence of carbonic anhydrase isoforms were observed by the SDS-PAGE technique [39]. The method was performed according to our previous studies [29,40,41]. After this procedure, a single band was seen for each isoform (Figure 1). Molecular weight of the enzyme was determined by using SynGene imaging tool (Figure 2).

Ammonium Sulfate Precipitation of PON1

20 ml of human serum was precipitated with solid ammonium sulfate at 60-80%, Centrifugation was carried out at 24000xg for 20 minutes during each precipitation. The pellet was dissolved in 100 mM Na-phosphate buffer pH 7.0 and dialyzed against same buffer [28].

Measurement of PON1 Activity

hPON1 activity was measured using paraoxon (diethyl p-nitrophenyl phosphate) as substrate (1 mM) in 50 mM glycine/NaOH (pH 10.5) including 1 mM CaCl₂. hPON1 assay was based on the measurement of p-nitrophenol at 412 nm [42].

RESULTS and DISCUSSION

Plants are employed for the treatment of many diseases date back to prehistory and all people have this ancient tradition. Due to their antioxidant, antimicrobial, antifungal antitumoral and anti-inflammatory activities, natural products have been used for a long time. Nowadays, plant extracts are consumed as health preservative materials among the people. This rate is almost over 80% of the world's population [43]. Most of the plants are rich in terms of phenolic compounds, benzophenones, xanthenes, bioflavonoid, flavonoid, terpenes as well as some metabolites such as anthraquinones, cyanates, oxalate, tannins and saponins [44]. Supportive roles of these compounds are indisputable in the living metabolism.

In this study, ethanol, methanol and water extractions of *Alcea rosea*, *Foeniculum vulgare*, *Elettaria cardamomum*,

Laurus azorica and *Lavandula stoechas* were prepared and measured effects of these samples against purified hPON1, hCA I and hCA II. For this reason, cytoplasmic CA isoforms, hCA I and hCA II, were purified from human erythrocytes with a simple and one step affinity method. Throughout the purification steps, CA-I was obtained with a specific activity of 768.5 U/mg proteins, with a yield of 62.25%. CA-II was also obtained with a specific activity of 930.34 U/mg proteins, with a yield of 23.27%. It is shown in Table 1.

The purity and molecular weight of the enzymes were determined approximately 29 kDa using the SDS-PAGE. (Figure 1 and Figure 2).

PON1 enzyme was obtained from human serum by only ammonium sulfate precipitation. Subsequently, ethanol, methanol and water extractions of the plant extracts were tested on purified enzyme activities. Almost all extracts inhibited the enzymes. It is important, because

both CAs and PON1 enzyme have esterase activity and the same reaction product is obtained at the end of the reaction (Figure 3).

Alcea rosea (*A.rosea*) is used as herbal remedy in folk medicine for treatment of different diseases such as inflammation of the kidneys and the uterus, kidney and urinary tract infections, malaria, rheumatism [45]. Some pharmacological effects including antibacterial, analgesic, anti-inflammatory and cytotoxic activities have been reported [46]. The inhibition results are expressed as IC_{50} (mg/mL) in the present study. As appeared from Table 1, hCAI, II and hPON1 activities are inhibited by *Alcea rosea* extracts. Only water extract did not have any inhibition or activation effect. Methanol extract was most effective for both hCAI and II. On the other hand, ethanol extract of *Alcea rosea* inhibited the hPON1 enzyme activity, strongly (0.074 mg/mL) (Table 2). There are some researches about the inhibition effects of *Alcea rosea* extracts on various

Table 1. Purification steps of hCA isoforms from human erythrocytes.

Step	Activity (EU/mg)	Protein (mg/mL)	Volume (mL)	Total Activity (EU)	Total Protein (mg)	Specific Activity (EU/mg)	Recovery (%)	Purification Fold
Hemolysate	130.0	4.60	38	4940.0	174.8	28.2	100	1
Sepharose-4B-L tyrosine-sulfanilamide affinity chromatography and dialysis for hCAI	416.6	0.542	7.5	3124.5	4.06	768.5	63.25	27.25
Sepharose-4B-L tyrosine-sulfanilamide affinity chromatography and dialysis for hCAII	383.3	0.412	3.0	1149.9	1.236	930.34	23.27	32.99



Figure 1. Sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) analysis of purified hCA isoforms. Lane 1: standard proteins, Lane 2: purified hCA-I, Lane 3: purified hCA-II.




Figure 2. Molecular weight of the hCA I and hCA II isozymes were calculated about 29 kDa by using SynGene imaging tool. Track 1 is the peaks of standard proteins. Track 2 is CA I's peak, and Track 3 is CA II's peak.




Figure 3. A) Mechanism of paraoxonase activity assay B) Mechanism of carbonic anhydrase esterase activity assay.

Table 2. Inhibitory effects of *Alcea rosea*, *Foeniculum vulgare*, *Elettaria cardamomum*, *Laurus azorica* and *Lavandula stoechas* different extracts against hCA I, hCA II and hPON1.

Plant Extracts	Solvent	hCA I IC ₅₀ (mg/mL)	hCA II IC ₅₀ (mg/mL)	hPON1 IC ₅₀ (mg/mL)
<i>Alcea rosea</i>	Water	---	1.270	1.721
	Ethanol	0.130	0.119	0.046
	Methanol	0.074	0.077	0.295
<i>Foeniculum vulgare</i>	Water	0.301	---	1.60
	Ethanol	---	0.162	0.073
	Methanol	0.183	0.129	0.200
<i>Elettaria cardamomum</i>	Water	0.991	1.400	1.361
	Ethanol	0.065	0.129	0.054
	Methanol	0.032	0.061	0.094
<i>Laurus azorica</i>	Water	0.842	1.170	0.732
	Ethanol	0.692	0.785	0.060
	Methanol	0.458	0.472	0.080
<i>Lavandula stoechas</i>	Water	0.119	0.330	---
	Ethanol	0.077	0.067	0.047
	Methanol	0.080	0.054	0.092

enzymes. For instance, Namjoyan et al. (2015) investigated the effects on diphenolase activity of mushroom tyrosinase of some plant extracts including *Alcea rosea* [47]. They found the IC_{50} value for *Alcea rosea* as 2.82 mg/mL.

Foeniculum vulgare is widespread in northern Anatolia. While the water extract did not have any effect on hCA II isozyme, ethanol extract did not have any effect on hCA I isozyme. Interestingly, hPON1 which is HDL-related antioxidant enzyme having an important role in the prevention of cardiovascular, was inhibited by each extract, and particularly, the enzyme activity was more decreased via ethanol extract (IC_{50} 0.073 mg/mL) Results were shown in Table 2.

Indeed, it is known that *F. vulgare* is a significant source of phenolic acids including of 1,4-O-di-caffeoylquinic acid, 1,3-O-di-caffeoylquinic acid, 1,5-O-di-caffeoylquinic acid, 3-O-caffeoylquinic acid, 5-O-caffeoylquinic acid, 4-O-caffeoylquinic acid, the flavonoids like rosmarinic, quercetin-3-rutinoside and eriodictyol-7-rutinoside [48]. They are phenolic compounds as well as antioxidant properties. For all this, we estimate that in our study, other factors in the content of the plant extract may be effective for inhibition of PON1 enzyme apart from phenolic compounds. In another study, thirteen compounds were isolated from a methanol extract of *F. vulgare* and tested for their inhibition on CYP3A4. 5-Methoxypsoralen (5-MOP) was seen to have strongest inhibition with IC_{50} value of 18.3 μ M and a mixed type of inhibition [49].

Elettaria cardamomum (L.), cardamom, is known as small cardamom and discriminated from large cardamom (*Amonum subulatum* Roxb.). Its oil has been employed in a wide range of beauty products [50]. Water, ethanol and methanol extracts of *Elettaria cardamomum* (L.) inhibited all enzymes. However, IC_{50} values of methanol extract were lower than other extract for hCA I and II isozymes. On the other hand, ethanol extract was seen to stronger inhibit the PON1 enzyme than other ones (Table 2; Figure 4).

Also, while IC_{50} values of hCA I and II obtained for methanol extracts of *Laurus azorica* was low, the highest inhibition for hPON1 was in ethanol extract (Table 2). The results were similar to *Laurus azorica* for *Lavandula stoechas* extracts (Table 2). There are some studies about the effects on enzymes of *L. Stoechas* extract. For example, Carrasco et al. (2015) investigated *L. Stoechas* extract as an enzyme target [51]. They prepared three samples of *L. Stoechas* and studied the effects of samples on lipoxygenase (LOX) activity. They determined the inhibition for LOX activity, significantly.

CONCLUSION

Carbonic anhydrase I and II isozymes were purified from human erythrocytes at one step by simple affinity technique. Paraoxonase 1 enzyme was obtained from human serum with partial purification using ammonium sulfate precipitation. Subsequently, water, ethanol and methanol extracts of *Alcea rosea*, *Foeniculum vulgare*, *Elettaria cardamomum*, *Laurus azorica* and *Lavandula*




Figure 4. IC_{50} graphs of the methanolic extracts of *E. cardamomum*. A) For carbonic anhydrase I B) For carbonic anhydrase II C) For paraoxonase enzyme activity.

stoechas were tested on hCA I, II and PON1 enzymes. IC₅₀ values were determined as mg/ml for each extract. According to the results, highest inhibition values were obtained in particular ethanol and methanol extracts of *Elettaria cardamomum*. As known, inhibition of the enzymes was crucial in the treatment of many diseases. However, using of the various plant extracts may lead to the formation some aggregates in the metabolism. Therefore, the next step of the study should be to determine the inhibitors containing extracts.

Acknowledgments

The authors declare that they have no conflict of interests.

References

- C.H. Chan, G.C. Ngoh, R. Yusoff, A brief review on anti diabetic plants: global distribution, active ingredients, extraction techniques and acting mechanisms, *Pharmacogn. Rev.*, 6 (2012) 22-28.
- N.J. Sucher, M.C. Carles, A pharmacological basis of herbal medicines for epilepsy, *Epilepsy Behav.*, 52 (2015) 308-318.
- P.R. Bhandari, A comment on effect of plant extracts on alzheimer's disease: an insight into therapeutic avenues, *J. Neurosci. Rural Pract.*, 4 (2013) 236-237.
- H.R. Vasanthi, N. Shrishrimal, D.K. Das, Phytochemicals from plants to combat cardiovascular disease, *Curr. Med. Chem.*, 19 (2012) 2242-2251.
- D. Krishnaiah, R. Sarbatly, R.Nithyanandam, A review of the antioxidant potential of medicinal plant species, *Food Bioprod. Process.*, 89 (2011) 217-233.
- A.E. Al-Snafi. The Pharmaceutical Importance of althaea officinalis and *Althaea rosea*: a review, *Int. J. Pharm. Tech. Res.*, 5 (2013) 1385-1387.
- P. Sadighara, S. Gharibi, A.M. Jafari, G.J. Khaniki, S. Salari, The antioxidant and flavonoids contents of althaea officinalis l. flowers based on their color, *Avicenna J. Phytomed.*, 2 (2012) 113-117.
- M. Kwiatkowska, D. Stepinski, K. Poplonska, A. Wojtczak, T.J. Polit, 'Elaioplasts' identified as lipotubuloids in *Althaea rosea*, *funkia sieboldiana* and *vanilla planifolia* contain lipid bodies connected with microtubules, *Acta Soc. Bot. Pol.*, 80 (2011) 211-219.
- R. Piccaglia, M. Marotti, Characterization of some Italian types of wild fennel (*Foeniculum Vulgare* Mill.), *J. Agric. Food Chem.*, 49 (2001) 239-44.
- S.B. Badgujar, V.V. Patel, A.H. Bandivdekar, *Foeniculum Vulgare* Mill: A review of its botany, phytochemistry, pharmacology, contemporary application, and toxicology, *Biomed. Res. Int.*, (2014) Article ID 842674.
- M.E. Leblebici, S. Machmudah, M. Sasaki, M. Goto, Antiradical efficiency of essential oils from plant seeds obtained by supercritical CO₂, soxhlet extraction, and hydrodistillation, *Sep. Sci. Technol.*, 48 (2012) 328-337.
- H. Al-Zuhair, B. El-Sayeh, HA. Ameen, H. Al-Shoora. Pharmacological studies of cardamom oil in animals, *Pharmacol. Res.*, 34, (1996) 79-82.
- I. Kubo, M. Himejima, H. Muroi, Antimicrobial activity of flavor components of cardamom *Elettaria cardamomum* (zingiberaceae) seed, *J. Agric Food Chem.*, 39, (1991) 1984-1986.
- O. Ouchikh, T. Chahed, R. Ksouri, M.B.Taarit, H. Faleh, C. Abdelly, M.E. Kchouk, B. Marzouk, The effects of extraction method on the measured tocopherol level and antioxidant activity of *L. nobilis* vegetative organs, *J. Food Comp. Anal.*, 24 (2011) 103-110.
- C. Da Porto, D. Decorti, Analysis of the volatile compounds of flowers and essential oils from *Lavandula angustifolia* cultivated in northeastern italy by headspace solid-phase microextraction coupled to gas chromatography-mass spectrometry, *Planta Med.*, 74 (2008) 182-187.
- A. Goren, G. Topçu, G. Bilsel, M. Bilsel, Z. Aydoğmuş, J.M. Pezzuto, The chemical constituents and biological activity of essential oil of *Lavandula stoechas* ssp. *Stoechas*, *Z. Naturforsch C*, 57 (2002) 797-800.
- T. Benabdelkader, A. Zitouni, Y. Guitton, F. Jullien, D. Maitre, H. Casabianca, L. Legendre, A. Kameli, Essential oils from wild populations of algerian *Lavandula stoechas* L.: composition, chemical variability, and in vitro biological properties, *Chem. Biodivers.*, 8 (2011) 937-953.
- D. Kaya, Ml. Alpaslan, E.S. Giray, S. Kirici, Diurnal, Ontogenetic and morphogenetic variability of *Lavandula stoechas* L. Ssp. *Stoechas* in East Mediterranean region, *Rev. Chim. (Bucharest Rom)*, 63 (2012) 749-753.
- M. Zuzarte, Gonçalves M.J, Cavaleiro C, Cruz M.T, Benzarti A, Marongiu B, Maxia A, Piras A, Salgueiro L. Antifungal and anti-inflammatory potential of *Lavandula stoechas* and thymus herba-barona essential oils, *Ind. Crops Prod.*, 44 (2013) 97-103.
- H.A. Aslan, Y. Demir, M.S. Özasan, F. Türkan, Ş. Beydemir, O.I. Kufrevioglu, The Behaviour of some chalcones on acetylcholinesterase and carbonic anhydrase activity, *Drug Chem. Toxicol.*, 4 (2018) 1-7.
- S. Mert, Z. Alim, MM. Işgor, S. Beydemir, R. Kasimogullari, The synthesis of novel pyrazole-3,4-dicarboxamides bearing 5-amino-1,3,4-thiadiazole-2-sulfonamide moiety with effective inhibitory activity against the isoforms of human cytosolic carbonic anhydrase I and II, *Bioorg. Chem.*, 68, (2016) 64-71.
- A. Topal, M. Atamanalp, E. Oruc, Y. Demir, S. Beydemir, A. Isik, In vivo changes in carbonic anhydrase activity and histopathology of gill and liver tissues after acute exposure to chlorpyrifos in rainbow trout, *Arh. Hig. Rada. Toksikol.*, 65, (2014) 377-385.
- Y. Demir, E. Oruç, A. Topal. Carbonic anhydrase activity responses and histopathological changes in gill and liver tissues after acute exposure to chromium in brown trout juveniles, *Hacettepe J. Biol. & Chem.*, 44 (2016) 515-523.
- A. Innocenti, I. Gulcin, A. Scozzafava, C.T. Supuran, Carbonic anhydrase inhibitors. antioxidant polyphenols effectively inhibit mammalian isoforms I-XV, *Bioorg. Med. Chem. Lett.*, 20 (2010) 5050-5053.
- H. Balseven, M.M. İşgor, S. Mert, Z. Alim, Ş. Beydemir, S. Ok, R. Kasimogullari, Facile synthesis and characterization of novel pyrazole-sulfonamides and their inhibition effects on human carbonic anhydrase isoenzymes, *Bioorg. Med. Chem.*, 21 (2013) 21-27.
- Z. Alim, N. Kilinc, M.M. Işgor, B. Sengul, S. Beydemir, Some anti-inflammatory agents inhibit esterase activities of human carbonic anhydrase isoforms I and II: an in vitro study, *Chem. Biol. Drug. Des.*, 86 (2015) 857-863.

27. H.A. Alici, D. Ekinci, Ş. Beydemir, Intravenous anesthetics inhibit human paraoxonase-1 (PON1) activity in vitro and in vivo, *Clin. Biochem.*, 41 (2008) 1384-1390.
28. S. Beydemir, Y. Demir, Antiepileptic drugs: Impacts on human serum paraoxonase-1, *J. Biochem. Mol. Toxicol.*, 31 (2017) 1-6.
29. Y. Demir, S. Beydemir, Purification, refolding, and characterization of recombinant human paraoxonase-1, *Turk J. Chem.*, 39 (2015) 764-776.
30. M. Oktay, I. Gülçin, O.I. Küfrevioğlu, Determination of in vitro antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts, *LWT- Food Sci. Technol.* 36 (2003) 263-271.
31. E.D. Kaya, H. Soyut, Ş. Beydemir, The toxicological impacts of some heavy metals on carbonic anhydrase from gilthead sea bream (*Sparus aurata*) gills, *Environ. Toxicol. Pharmacol.*, 39 (2015) 825-832.
32. D. Ekinci, Ş. Beydemir, Z. Alim, Some drugs inhibit in vitro hydratase and esterase activities of human carbonic anhydrase-I and II, *Pharmacol. Rep.*, 59 (2007) 580-587.
33. Z. Alim, B. Çamur, Ş. Beydemir, O.I. Küfrevioğlu, The correlation between some metal concentrations and carbonic anhydrase activity in tuna (*thunnus thynnus linnaeus*, 1758) gill, *Hacettepe J. Biol. & Chem.*, 42 (2014) 219 – 224.
34. J.A. Verpoorte, S. Mehta, J.T. Edsall, Esterase activities of human carbonic anhydrases B and C, *J. Biol. Chem.*, 242 (1967) 4221-4229.
35. M.M. Bradford, Rapid and sensitive method for quantitation of microgram quantities of protein utilizing principle of protein-dye binding, *Anal. Biochem.*, 72 (1976) 248-254.
36. Y. Demir, M. Isik, I. Gulcin, Ş. Beydemir, Phenolic compounds inhibit the aldose reductase enzyme from the sheep kidney, *J. Biochem. Mol. Toxicol.*, 31 (2017) e21935.
37. Y. Demir, M. Şenol Kotan, N. Dikbaş, Ş. Beydemir, Phytase from *Weissella halotolerans*: purification, partial characterisation and the effect of some metals, *Int. J. Food Prop.*, 20 (2017) 2127-2137.
38. M.S. Ozaslan, Y. Demir, O.I. Kufrevioğlu, M. Ciftci, Some metals inhibit the glutathione S-transferase from Van Lake fish gills, *J. Biochem. Mol. Toxicol.*, 31 (2017) e21967.
39. U.K. Laemmli, Cleavage of structural proteins during assembly of head of bacteriophage-T4, *Nature*, 227 (1970) 680–683.
40. M. Kirici, M. Kirici, Y. Demir, Ş. Beydemir, M. Atamanalp, The effect of Al³⁺ and Hg²⁺ on glucose 6-phosphate dehydrogenase from capoetaumbra kidney, *Appl. Ecol. Env. Res.*, 14 (2016) 253-264
41. Y. Demir, B. Şengül, B. Ergun, Ş. Beydemir, Alcohol dehydrogenase from sheep liver: Purification, characterization and impacts of some antibiotics, *Iğdır Univ. J. Inst. Sci. & Tech.*, 7 (2017) 151-159.
42. M. Isik, Y. Demir, M. Kirici, R. Demir, F. Simsek, Ş. Beydemir, Changes in the anti-oxidant system in adult epilepsy patients receiving anti-epileptic drugs, *Arch. Physiol. Biochem.*, 121 (2015) 97-102.
43. H. Sahin, Z. Can, O. Yildiz, S. Kolayli, A.G. Innocenti, Scozzafava, C.T. Supuran, Inhibition of carbonic anhydrase isozymes i and ii with natural products extracted from plants, mushrooms and honey, *J. Enzyme. Inhib. Med. Chem.*, 27 (2012) 395-402.
44. M.F. Asaolu, Chemical compositions and phytochemical screening of the seeds of garcinia kola (bitter kola), *Pak. J. Sci. Ind. Res.*, 46 (2003) 145-147.
45. M. Munir, A. Hussain, I. Ul-Haq, R. Qureshi, M. Munazir, M. Rshad, M. Khan, Callogenesis potential of cotyledonary explants of *Althaea rosea* from Pakistan, *Pak. J. Bot.*, 44 (2012) 271-275.
46. T. Mert, T. Erdogan, B. Kivcak, T. Ozturk, Antimicrobial and cytotoxic activities of the extracts obtained from the flowers of *Alcea rosea* L., *J. HUJPHARM*, 30 (2010) 17-24.
47. F. Namjoyan, A. Jahangiri, M.E. Azemi, E. Arkian, H. Mousavi, Inhibitory effects of physalis alkekengi l., *Alcea rosea* l., *Bunium persicum* b. fedtsch. and *Marrubium vulgare* l. on mushroom tyrosinase, *Jundishapur. J. Nat. Pharm. Prod.*, 20 (2015) 23356.
48. M. Faudale, F. Viladomat, J. Bastida, F. Poli, C. Codina, Antioxidant activity and phenolic composition of wild, edible, and medicinal fennel from different mediterranean countries, *J. Agric. Food Chem.*, 56 (2008) 1912-1920.
49. S.F. Subehan Zaidi, S. Kadota, Y. Tezuka, Inhibition on human liver cytochrome p450 3a4 by constituents of fennel (*Foeniculum vulgare*): identification and characterization of a mechanism-based inactivator, *J. Agric. Food Chem.*, 55 (2007) 10162-67.
50. B. Marongiu, A. Piras, S. Porcedda, Comparative analysis of the oil and supercritical CO₂ extract of *Elettaria Cardamomum* (L.) Maton, *J. Agric. Food Chem.*, 52 (2004) 6278-682.
51. A. Carrasco, O.R. Vanessa, M.G. Ramiro, T. Virginia, J. Tudela *Lavandula stoechas* essential oil from spain: aromatic profile determined by gas chromatography–mass spectrometry, antioxidant and lipoxigenase inhibitory bioactivities, *Ind. Crops Prod.*, 73 (2015) 16-27.